Department of Mathematics

Preprint MPS_2010-08

20 March 2010

Eigenvalue problem meets Sierpinski triangle: computing the spectrum of a nonself-adjoint random operator

by

Simon N. Chandler-Wilde, Ratchanikorn Chonchaiya and Marko Lindner



Eigenvalue problem meets Sierpinski triangle: computing the spectrum of a non-self-adjoint random operator

Simon N. Chandler-Wilde, Ratchanikorn Chonchaiya and Marko Lindner

March 20, 2010

Abstract. The purpose of this paper is to prove that the spectrum of the non-self-adjoint one-particle Hamiltonian proposed by J. Feinberg and A. Zee (Phys. Rev. E 59 (1999), 6433(6443) has interior points. We do this by rst recalling that the spectrum of this random operator is the union of the set of ` $^{\infty}$ eigenvalues of all in nite matrices with the same structure. We then construct an in nite matrix of this structure for which every point of the open unit disk is an ` $^{\infty}$ eigenvalue, this following from the fact that the components of the eigenvector are polynomials in the spectral parameter whose non-zero coe cients are ±1's, forming the pattern of an in nite discrete Sierpinski triangle.

Mathematics subject classi cation (2000): Primary 47B80; Secondary 47A10, 47B36. *Keywords:* random matrix, spectral theory, Jacobi matrix, disordered systems.

1 Introduction and Notations

In this paper we study in nite matrices of the form



(1)

with $b_k 2f \ 1g := f \ 1; +1g$ for all $k \ 2Z$

Physicists have studied the operator A^b as the (non-self-adjoint) Hamiltonian of a particle hopping (asymmetrically) on a 1-dimensional lattice [15, 16, 9, 22]. Applications of such and related Hamiltonians, especially examples with random diagonals, include vortex line pinning in superconductors and growth models in population biology. The particular model (1) was proposed by Feinberg and Zee in [15], and some properties of its spectrum have been studied in [9, 22] (also see Paragraph 37, in particular Figure 37.7c, in [38]).

In all these studies the focus is on the case of a random sequence $b \ 2 \ f \ 1g^Z$. A related but completely deterministic concept is that of a pseudo-ergodic sequence. In accordance with Davies [11], we call $b \ 2 \ f \ 1g^Z$ pseudo-ergodic if every nite pattern of 1's can be found somewhere (as a string of consecutive entries) in *b*. If *b* is pseudo-ergodic (which is almost surely the case if all b_k , $k \ 2 \ Z$, are independent (or at least not fully correlated) samples from a random variable with values in *f* 1*g* and nonzero probability for both +1 and 1) then, as a consequence of [7] (also see [6, 8, 29, 30] and cf. [11]), it holds that

spec
$$A^b$$
 = spec_{ess} A^b = $\begin{bmatrix} spec A^c = \\ c2f \ 1g^{\mathbb{Z}} \\ c2f \ 1g^{\mathbb{Z}} \\ c2f \ 1g^{\mathbb{Z}} \end{bmatrix}$ (2)

The contribution of [7] is the third $\=$ " sign in (2) that enables, or at least simpli es, the explicit computation of the spectra of particular pseudo-ergodic operators in [6, 8, 29]. The rst $\=$ " sign in (2) follows immediately from the second; the second comes from the Fredholm theory of much more general operators and is typically expressed in the language of so-called limit operators [34, 35, 27, 8]. (A similar equality, often with the closure taken of the union of spectra, can be found in the literature on spectral properties of Schrodinger and more general Jacobi operators [32, 4, 10, 11, 21, 1, 31, 17, 18, 19, 20, 33, 26, 25, 36, 37]. The three last papers also shed some light on the role of limit operators in the study of the absolutely continuous spectrum.)

Note that, by (2), the spectrum of A^b does not depend on the actual sequence b { as long as it is pseudo-ergodic. In [6] we obtain information about the spectrum, pseudospectrum and numerical range of the bi-in nite matrix operator A^b , its contraction A^b_+ to the positive half axis (a semi-in nite matrix) and the nite sections A^b_n which, for $n \ge N$, are n = n submatrices of (1). Explicitly and precisely, these related matrices are

where in the case n = 1 we set $A_1^b = (0)$. We explore in some detail in [6] the interrelations between the spectra and pseudospectra of A^b , A_+^b and A_n^b . Here, for " > 0 and a bounder A^b and A_n^b . Here, for " > 0 and a bounder A^b and A_n^b . **b)** Provided the \positive" part of the sequence b (by which we mean $(b_k)_{k \ge N}$) is itself pseudoergodic (contains every nite pattern of 1's), then, for all " 0 one has

 $\operatorname{spec}_{"} A^{b} = \operatorname{spec}_{"} A^{b}_{+}$:

c) l^{b}_{A} calgleq of

literature) whether spec A^b has positive Lebesgue measure, in particular whether it has interior points. Related to this question, Holz et al. [22, Sections I, V, VI], conjecture that clos ($_1$) spec A^b has a fractal dimension in the range (1;2), and so has zero Lebesgue measure.

The purpose of the current paper is to shed light on these questions by constructing a sequence $c \ 2 \ f \ 1g^{Z}$ for which spec¹_{point} A^{c} contains the open unit disk. As a consequence of formula (2) and the closedness of spectra, this shows that spec A^{b} contains the closed unit disk and therefore has dimension 2 and a positive Lebesgue measure. This is the main result of the next section. Intriguingly we will see that the sequence constructed, while rather irregular, is such that each in the unit disk is an eigenvalue of A^{c} in 1 (Z), and with a vector whose components are polynomials in with coe cients forming the regular self-similar pattern of a discrete Sierpinski triangle.

We will nish the paper with our own conjecture on the geometry of clos (1) and spec A^b .



Figure 1.2: Our gure shows the sets n := n, 0 of all $n \times n$ matrix eigenvalues, as defined in (4), for n = 1, ..., 30. Note that in the first pictures (with only a few eigenvalues), we have used heavier pixels for the sake of visibility. By (5), each of the sets with n = 1, 2, ..., 14 in this gure is contained, respectively, in the set number 2n + 2 of Figure 1.1.



Figure 1.3: This is a zoom into $_{25}$ { the 25th picture of Figure 1.2. The location of this zoom is near the point 1 + *i*, which is the midpoint of the northeast edge of the square clos ($W(A^b)$) = conv{2; -2; 2i; -2i}. The picture clearly suggests self-similar features of the set $_{25}$.

2 A sequence c for which spec A^c contains the unit disk

The formula (2) for the spectrum of A^b when $b \ 2f \ 1g^Z$ is pseudo-ergodic motivates the following approach to decide whether a given point $2 \ C$ is in spec A^b or not: look for a sequence $c \ 2f \ 1g^Z$ such that $2 \ \text{spec}_{\text{point}}^1 A^c$, in other words, such that there exists a non-zero $u \ 2^{1} \ (Z)$ with

 $A^{c}u = u$, i.e.

$$U_{i+1} = U_i \qquad C_i U_{i-1}$$
 (6)

for $i 2 \mathbb{Z}$. If such a sequence *c* exists then 2

and so on. Explicitly, it is easy to check that, for i = 3, the solution of (6) with initial conditions $u_0 = 0$ and $u_1 = 1$ is given by the characteristic polynomial



Thus, for *i* 3, u_i is a polynomial of degree *i* 1 in with coe cients depending on c_2 ; ...; c_{i-1} . We will aim to achieve that *u* be a bounded sequence at least for *j j* < 1. With this in mind we should try to keep the coe cients of these polynomials small. Precisely, our strategy will be to try to choose c_1 ; c_2 ; ...; 2 *f* 1*g* such that each u_i is a polynomial in with coe cients in *f* 1;0;1*g*. The following table, where we abbreviate 1 by , +1 by +, and 0 by a space, suggests that this seems to be possible.



For $i; j \ge N$, denote the coe cient of j = 1 in the polynomial u = 0 9.464 / SQq1 87038 - .4904 Td [iN

1

Proposition 2.1 De ne the sequence $c \ 2 \ f \ 1g^Z$, for positive indices by $c_1 = 1$ and by the requirement that

 $c_{2i} = c_{2i-1}c_i$ and $c_{2i+1} = c_{2i};$ i = 1/2; ...;

Now suppose i + j is odd. Then, by (10) and the inductive hypothesis,

$$\begin{array}{rcl} p_{2i & 1;2j & 1} & = & p_{2i & 2;2j & 2} & C_{2i & 2} & p_{2i & 3;2j & 1} \\ & & = & 0 & C_{2i & 3} & C_{i & 1} & p_{i & 1;j} & = & C_{2i & 1} & p_{i & 1;j} \end{array}$$

since $c_{2i} = c_{2i} = c_{2i} c_{i} c_{i-1}$. By (10) and the inductive hypothesis and noting that i + j is even,

$$\begin{array}{rcl} \rho_{2i;2j} &=& \rho_{2i-1;2j-1} & C_{2i-1} \rho_{2i-2;2j} \\ &=& C_{2i-1} \rho_{i-1;j} & C_{2i-1} \rho_{i-1;j} &=& 0; \end{array}$$

This completes the proof of (iv), and (v) follows from (iv) by a simple induction argument.

To see that (vi) is true, observe rst that, from (i), (iii), and (iv) (and cf. Remark 2.2), it holds for $i^{0}; j^{0} \ge N$ that $(i^{0}; j^{0}) \ge S$ i , for some $i; j \ge N$ either (a) $(i^{0}; j^{0}) = (2i; 2j)$ and $(i; j) \ge S$; or (b) $(i^{0}; j^{0}) = (2i - 1; 2j - 1)$ and $(i; j) \ge S$ or $(i - 1; j) \ge S$. From this it follows that S = T(S).

De ne a metric *d* on by

$$d(;) := \begin{array}{c} \times \\ 2^{i j}; ; 2 : \\ (i;j)2([)n(\backslash) \end{array}$$

Then

if $(1/1) \mathcal{B}([]) n([])$. Let $_1 := f 2 : (1/1) 2 g$. Then $T(_1) = _1$ and, by (14), T is a contraction mapping on $_1$. Thus, by the contraction mapping theorem, T has a unique xed point in $_1$, which is the set S, and, if $_1 2 _1$ and $_{n+1} := T(_n)$, $n 2 \mathbb{N}$, then $d(_n; S) ! 0$ as n ! 1. In particular, $d(S_n; S) ! 0$ as n ! 1. Since also (by an easy induction argument) $S_1 = S_2 = ...$, it follows that $S = [_{n 2 \mathbb{N}} S_n$.

Define v_i for i = 0, 1, ... by $v_i := d_i u_i$, which implies that $v_0 = 0$, and set $v_1 = 1$. Then, since u_i is defined uniquely for i = 0 by the requirement that it satisfy (6) for i = 0 with the initial conditions that $u_0 = 0$ and $u_1 = 1$, to show (*vii*) it is enough to check that the sequence v_i satisfies (6) for i = 0, i.e. that

 $V_{i+1} = V_i \quad C_i V_{i-1}; \quad i = 0; 1; ...;$

But $v_1 = v_0 + c_0 v_1 = 1 + c_0 d_1 u_1 = 0$, so the equation holds for i = 0, and, for $i \ge N$,

$$V_{i+1} V_{i} + C_{i}V_{i-1} = d_{i-1}U_{i-1} d_{i}U_{i} + C_{i+1}d_{i+1}U_{i+1}$$

= $(d_{i-1} C_{i}C_{i+1}d_{i+1})U_{i-1} (d_{i-1} C_{i+1}d_{i+1})U_{i}$

since $u_{i+1} = u_i$ $c_i u_{i-1}$. Since $u_0 = 0$, the right hand side of this last equation is zero for $i \ge 0$ provided that $d_i = c_{i+1}d_{i+1}$ for $i \ge 0$. But this follows from the denitions of the sequences c and d.

Remark 2.4 The standard in nite discrete Sierpinski triangle (e.g. [24]) is the set $S = \sum_{n \ge N} S_n$, where $S_1 := f(1;1)g$ and the sets S_n , n = 2;3;..., are defined ned recursively by $S_{n+1} := 2S_n + V$, where V := f(0;0); (-1; -1); (0; -1)g. One instance where S arises is as the pattern of odd coe cients in Pascal's triangle: for $i \ge N$ and j = 1;...;i the coe cient of x^{j-1} in $(1 + x)^{j-1}$ is odd i $(i;j) \ge S$, so that the discrete Sierpinski triangle is often referred to as Pascal's triangle modulo 2 (e.g. [13]). Proposition 2.1(vi) (cf. Remark 2.2) makes clear that the pattern $S = N^2$ of the non-zero coe cients in table (7) is essentially that of the standard discrete Sierpinski triangle S; indeed, the sets S and S are connected by a linear mapping: $(i;j) \ge S$ i $(2i = j;j) \ge S$, for $i;j \ge N$. \Box

Remark 2.5 Note that the sequence *c* from Proposition 2.1 is not pseudo-ergodic since, by $c_{2i+1} = c_{2i}$, the patterns $\backslash + + + "$ and \backslash " can never occur as consecutive entries in the sequence *c*. \Box

Based on Theorems 1.1 and 2.3 and the numerical results displayed in Figures 1.1 and 1.2, we make the following conjecture.

.68.915238 Tf 5A 0 T CC (xing) = Cture 9. WAG 2 (nhje): ture f that T tus ((.)] T) = EcTos (W) 18. Sphel(()) k361(=)-360(e(wing)-334.311 0 Td [(.)] T J E238 Tf !